

ON THE OPTIMAL WEIGHT FUNCTION IN THE GOLDSTON-PINTZ-YILDIRIM METHOD FOR FINDING SMALL GAPS BETWEEN CONSECUTIVE PRIMES

JÁNOS PINTZ, SZILÁRD RÉVÉSZ, AND BÁLINT FARKAS

ABSTRACT. We work out the optimization problem, initiated by K. Soundararajan, for the choice of the underlying polynomial P used in the construction of the weight function in the Goldston–Pintz–Yıldırım method for finding small gaps between primes. First we reformulate to a maximization problem on $L^2[0, 1]$ for a self-adjoint operator T , the norm of which is then the maximal eigenvalue of T . To find eigenfunctions and eigenvalues, we derive a differential equation which can be explicitly solved. The aimed maximal value is $S(k) = 4/(k + ck^{1/3})$, achieved by the $k - 1^{\text{st}}$ integral of $x^{1-k/2}J_{k-2}(\alpha_1\sqrt{x})$, where $\alpha_1 \sim ck^{1/3}$ is the first positive root of the $k - 2^{\text{nd}}$ Bessel function J_{k-2} . As this naturally gives rise to a number of technical problems in the application of the GPY method, we also construct a polynomial P which is a simpler function yet it furnishes an approximately optimal extremal quantity, $4/(k + Ck^{1/3})$ with some other constant C . In the forthcoming paper of J. Pintz [8] it is indeed shown how this quasi-optimal choice of the polynomial in the weight finally can exploit the GPY method to its theoretical limits.

1. INTRODUCTION

1.1. The extremal problem as given by Soundararajan. In his work [11] Soundararajan presents and analyzes the proof of Goldston–Pintz–Yıldırım yielding small gaps between primes. Among others he raises and answers one of the most important problems of the field: Is it possible to modify the weight function $a(n)$ in such a way that the method would lead to infinitely many bounded gaps between consecutive primes. If we consider the weight functions in full generality, that is all functions $a(n)$, then this leads essentially to a tautology. For example, defining $a(n) = 1$ if both n and $n + 2$ are primes, and otherwise setting $a(n) = 0$, the summatory function of $a(n)$ describes the number of twin primes up to x . Thus we cannot hope an asymptotic evaluation of the summatory function. We briefly describe the feasible choices of the weight function a . Let us take an admissible k -tuple $\mathcal{H} = \{h_1, \dots, h_k\}$ meaning that there is no prime p with the property that the elements h_i of \mathcal{H} cover all residue classes mod p . Let $P_{\mathcal{H}}(n) = \prod_{i=1}^k (n + h_i)$ and let us define $\lambda_d = \mu(d)P(\frac{\log(R/d)}{\log R})$ with a nice function P , for example a polynomial, with the additional property $\lambda_1 = 1$, which is equivalent to $P(1) = 1$. Afterwards we reduce our choice of $a(n)$ to those of type $a(n) = \sum_{d \leq R, d|P_{\mathcal{H}}(n)} \lambda_d$ and try to evaluate the summatory function of $a(n)$ and that of $a(n)\chi(n + h)$, where h is an arbitrary number

Date: May 22, 2013.

2000 *Mathematics Subject Classification.* Primary 11N05, 47A75. Secondary 47A53, 49J05, 49K15, 49N10.

Key words and phrases. Prime distribution, gaps between primes, Goldston–Pintz–Yıldırım method, selfadjoint Fredholm type operator, Bessel differential equation, Bessel functions of the first and second kind.

Supported in part by the Hungarian National Foundation for Scientific Research, Project # K-81658, K-100291, K-100461 and NK-104183. Work done in the framework of the project ERC-AdG 228005.

with $h < \log n$ and χ is the characteristic function of the primes. (In case of bounded gaps between primes it is sufficient to consider the case when $h = h_i, i = 1, 2, \dots, k$.)

Soundararajan explains, how the optimal weight function $a(n)$, hence λ_d , should be chosen to obtain best result: see formula (8) in [11]. In order to get this optimum, he also explains the choice $\lambda_d := \mu(d)P\left(\frac{\log(R/d)}{\log R}\right)$ where P is some suitably nice function, like a polynomial or at least a sufficiently many times (at least k times) differentiable, smooth function on $[0, 1]$ (or at least on $[0, 1)$), *vanishing at least in the order k at 0*, and satisfying the normalization $P(1) = 1$. Then, according to the analysis by Soundararajan, the optimal choice for $a(n)$ and λ_d is equivalent to looking for the maximal possible value of (12) of [11], i.e., to determining

$$(1) \quad S(k) := \sup_P \left(\int_0^1 \frac{x^{k-2}}{(k-2)!} (P^{(k-1)}(1-x))^2 dx \right) / \left(\int_0^1 \frac{x^{k-1}}{(k-1)!} (P^{(k)}(1-x))^2 dx \right),$$

where the set of functions P , to be taken into account in the supremum, can be the set of certain polynomials as before, or more generally a family of functions subject to some conditions.

Soundararajan [11] shows that the question whether we are able to find in this way infinitely many bounded gaps between primes is equivalent to the problem whether there exists any natural number k with $S(k) > 4/k$. Then he mentions that the opposite inequality $S(k) < 4/k$ holds for all k and therefore the method cannot yield infinitely many bounded prime gaps. (In an earlier unpublished note [10] he gives the short proof of this fact; we will reproduce this in §2.2. His considerations also lead easily to the stronger inequality $S(k) < 4/(k + c \log k)$, cf. §2.2). Although his work answered negatively the above mentioned central problem, it gave some hints but did not answer the question: What is the best weight function that can be chosen, and what size of gaps are implied by it? In their work [5] Goldston, Pintz and Yıldırım showed that if one takes $P(x) = x^{k+\ell}$, where k and ℓ are allowed to tend to infinity with the size N of the primes considered, then with several essential modifications of the original method one can reach infinitely many prime gaps of size essentially $\sqrt{\log p}$. (To have an idea of the difficulties it is enough to mention that the rather condensed proof of the result needs about 40 additional pages beyond the original one, presented with many details and explanations in [4]. However, a shortened, simplified and more condensed version [6] needs only 5 pages). In this case $\ell = c\sqrt{k}$ and the value of the fraction (1) is $4/(k + c'\sqrt{k})$ for the given choice of $P(x) = x^{k+\ell}$. Beyond the mentioned important fact that k and ℓ are unbounded in [5], the scheme of the proof is similar but not the same as in the simplified version of Soundararajan [11]. However, a careful analysis suggests that in order to find the limits of the method it is necessary (but as discussed a little later, not necessarily sufficient) to find the size of $S(k)$ as k tends to infinity together with the function P which yields a maximum (if it exists) in the supremum, or at least a function P which yields a value “enough close” to the supremum.

1.2. Conditions and normalizations. Before proceeding, let us discuss right here the issue of conditions and normalizations in the formulation of this maximization problem. First, it is clear that $P^{(k)}$ remains unchanged, if we add any constant to $P^{(k-1)}$. Thus the extremal problem becomes unbounded under addition of a free constant, hence at least some conditions must certainly control this divergence.

In the number theory construction of Goldston–Pintz–Yıldırım, (by now generally abbreviated as the “GPY method”) the natural restriction is that P must be a polynomial divisible by x^k —or, if we try to generalize the method, then a k -times continuously differentiable function with $P^{(j)}$ vanishing at 0 for $j = 0, \dots, k - 1$. That is $P(x) = \frac{P^{(k)}(0)}{k!}x^k + o(x^{k+1})$. The reason for that is the fact that the whole idea hinges upon the use of the generalized Möbius inversion, more precisely of the Λ_j function, which must be zero for numbers having at least k prime factors—always satisfied by the numbers represented by the product form $(n + h_1) \cdots (n + h_k)$ in the construction. So for any meaningful weight function we need to use weights not containing any smaller power x^j than x^k . In other words, we should assume here P having a zero of order k , i.e., $P(0) = P'(0) = \dots = P^{(k-1)}(0) = 0$, while $P^{(k)}(0)$ can be arbitrary.

The analysis of Soundararajan exposed the question, whether a linear combination of monomials, i.e., a polynomial, or perhaps some more sophisticated choice of a weight function, may perhaps improve even upon this. We can say that the *theoretical limit of the GPY method* is the result, obtainable in principle by a choice of the weight function P maximizing the extremal quantity (1). Yet it is to be noted that the technicalities of GPY are far more substantial than to simply “substituting any P ” in it would automatically lead to a result—it is not even that clear, what result would follow from a given weight function. Therefore, to test the limits of the GPY method, we should break our approach into two parts. First, we look for the optimization of the weight P , in the sense of (1), and second, we extend the GPY method using that weight function. This paper is concerned with this first question, and the second part of this program is carried out in [8].

The aim of the present analysis is to settle the issue of optimization in Problem (1). We find the optimal order, and the maximizer of the problem (1), furthermore, as this maximum can be achieved by a relatively sophisticated choice of the weight function P —actually a transformed Bessel function—we also construct a polynomial weight which is approximately optimal in (1).

Part of these results were reached by J. B. Conrey and his colleagues at the American Institute of Mathematics already in 2005. Using a calculus of variation argument they found the Bessel function J_{k-2} and made some calculations for concrete values of k (without analyzing the case $k \rightarrow \infty$). The fact that the Bessel functions may perform better than polynomials in the GPY method is also briefly noted in the book of J. B. Friedlander and H. Iwaniec [3] without going into details.

1.3. Structure of the paper. In this paper we proceed along the following course.

Interpreting the problem in the widest possible function class which makes sense (i.e., when at least the occurring integrals exist finitely) in Section 3 we make several further reformulations until we arrive at a maximization problem in the Hilbert space $L^2[0, 1]$. Exploiting the rich structure of Hilbert spaces, and the particular properties of the reformulation as a certain quadratic form with a Fredholm-type operator, we derive existence of maximizing functions in this wide function class. Then we also exploit the concrete form of the kernel in our Fredholm-type operator and compute that the maximizers, or, more generally, eigenfunctions, are necessarily smooth. Furthermore, in Section 4 we find that these eigenfunctions satisfy certain differential equations. The solutions are then found to be transformed variants of certain Bessel functions. Also it turns out that the solutions are analytic, and they yield a function value in the extremal problem directly related to the choice of a parameter, which, due to the initial value restriction $P^{(k-1)}(0) = 0$, must be a zero of the arising Bessel function J_{k-2} . Finally these combine to the full description

of the maximal value $S(k)$ together with the precise form of the extremal function. From the well-known asymptotic formula for the first zero of the Bessel function J_m , when $m \rightarrow \infty$, we derive that $S(k)$ is precisely asymptotic to $\frac{4}{k+ck^{1/3}}$ with a concretely known constant $c = 3.7115\dots$

Unfortunately, in spite of analyticity and power series expansion, the found extremal function is too complicated to be used in the number theory method of GPY. Basically, we need restrictions on the degree and the coefficient size for the powers appearing in the weight function P to make the complicated method work in a technically feasible way. As discussed in Section 5, not even calculations using the power series expansion of Bessel functions lead to feasible expressions. Therefore, finally we look for quasi-optimal polynomials, which still achieve close to extremal values. The result of the last section is the concrete construction of a polynomial P satisfying the needed technical requirements and still achieving in (1) a ratio of the order $\frac{4}{k+ck^{1/3}}$ with some other constant C . That suffices in the method of GPY, because the value of the constant C does not increase the order, only the arising constants, in the final result.

Settling the issue of the search of optimal and quasi-optimal weights, the door opens up for revisiting the method of GPY and not only improve upon all the known results, but also push the available techniques to the theoretical limits of that method. This closely connected work is carried out in the paper [8].

2. REFORMULATIONS AND THE FINITENESS OF $S(k)$

2.1. Reformulations. The normalization $P(1) = 1$ is rather inconvenient because the next reformulation (still following Soundararajan) is to put $Q(y) := P^{(k-1)}(y)$, a completely logical step in view of the fact that no values of P , P' , etc. $P^{(k-2)}$ occur in the actual optimization problem (1) and that the still occurring $P^{(k-1)}$ and $P^{(k)}$ can be nicely expressed as Q and Q' . So in line with the restriction that P vanishes at least to the order k at 0, following Soundararajan we write

$$\begin{aligned} P(x) &= \int_0^x \int_0^{x_1} \dots \int_0^{x_{k-2}} P^{(k-1)}(x_{k-1}) dx_{k-1} \dots dx_1 = \int_0^x \int_0^{x_1} \dots \int_0^{x_{k-1}} P^{(k)}(x_k) dx_{k-1} \dots dx_1 \\ &= \int_0^x \int_0^{x_1} \dots \int_0^{x_{k-2}} Q(x_{k-1}) dx_{k-1} \dots dx_1 = \int_0^x \int_0^{x_1} \dots \int_0^{x_{k-1}} Q'(x_k) dx_k \dots dx_1. \end{aligned}$$

Therefore, $P(0) = \dots = P^{(k-2)}(0) = P^{(k-1)}(0) = 0$ transforms to the simpler requirement that $Q(0) = 0$, while the corresponding P is obtained by the above integrals directly. Let us record one more thing here: The condition that $P(1) = 1$, expressed in terms of Q , is a *linear restriction*, as $I(Q) := (P(1) =) \int_0^1 \int_0^{x_1} \dots \int_0^{x_{k-2}} Q(x_{k-1}) dx_{k-1} \dots dx_1$ is just a linear functional on the function Q . To express it in a more condensed, closed form, we may apply Fubini's theorem to get a representation in the form of the well-known Fubini integral

$$\begin{aligned} P(x) &= \int_0^x Q(t) \frac{(x-t)^{k-2}}{(k-2)!} dt, \\ P(1) &= \int_0^1 Q(t) \frac{(1-t)^{k-2}}{(k-2)!} dt = \int_0^1 Q(1-y) \frac{y^{k-2}}{(k-2)!} dy. \end{aligned}$$

Note the similarity to the numerator of the quotient in (1). It is thus immediate by the Cauchy-Schwarz inequality that $P(1)$ is a finite, convergent integral whenever the

Lebesgue integral $\int_0^1 Q^2(1-y) \frac{y^{k-2}}{(k-2)!} dy$ exists. That is, no special requirement is needed to this effect once we guarantee that the numerator and denominator in (1) are well-defined.

In all, we were to look for maximum in the family

$$(2) \quad \mathcal{P} := \left\{ P(x) = \int_0^x \int_0^{x_1} \dots \int_0^{x_{k-2}} P^{(k-1)}(x_{k-1}) dx_{k-1} \dots dx_1, \quad P(1) = 1, \quad P^{(k-1)}(0) = 0 \right\},$$

but following Soundararajan we changed the setup to

$$(3) \quad \mathcal{Q}_1 := \left\{ Q : \int_0^1 Q(1-y) \frac{y^{k-2}}{(k-2)!} dy = 1, \quad Q(0) = 0 \right\},$$

where now Q can be any (say, continuously differentiable) function satisfying the requirements. The quantity we seek to maximize is then expressed as

$$(4) \quad \frac{\int_0^1 ny^{n-1}Q^2(1-y)dy}{\int_0^1 y^n Q^2(1-y)dy} \quad (n := k-1),$$

which again is a fraction of two expressions, both *quadratic homogeneous* in Q . Therefore, the ratio will be the same for cQ with any $c \neq 0$ and the original question can thus be rewritten as looking for the supremum of these quantities among functions in

$$\mathcal{Q}^* := \left\{ Q : \int_0^1 y^{k-2} Q(1-y) dy \neq 0, \quad Q(0) = 0 \right\}.$$

Continuity of Q' is not indispensable, but of course the ratio must be a ratio of finite quantities, with a nonzero and finite denominator, hence we need still to restrict considerations to functions $Q' \not\equiv 0$ or, in general allowing discontinuous functions, Q' not zero almost everywhere and also satisfying $\int_0^1 x^n Q^2(1-x) dx < \infty$. This latter condition also implies (by a partial integration and the Cauchy–Schwarz inequality) that even $\int nx^{n-1}Q^2(1-x)dx < \infty$, as needed. Furthermore, together with the restriction that $Q(0) = 0$, we see that Q is constant if only $Q \equiv 0$, so we need to exclude only this obviously singular case, and otherwise with the condition $Q(0) = 0$ a finite, fixed bound can be obtained for the ratio in question, hence $S(k)$ exists finitely at least as a supremum.

Let us observe that the condition that $P(1) \neq 0$, is a condition that a linear functional on the function space of admissible functions Q should not vanish. In other words, the subset which falls out of consideration for not meeting this condition is the kernel subspace of the linear functional, which is—in view of the fact that the functional itself is not identically zero (e.g., look at nonnegative, not identically zero functions Q which certainly provide nonzero functional values)—of codimension one, that is a hyperplane \mathcal{H} of our linear function space \mathcal{X} (whatever choice of the function space and respective norm we make later on). Therefore, \mathcal{H} is contained in the closure of $\mathcal{X} \setminus \mathcal{H}$ and thus—by continuity¹ of the given ratio which we maximize—the supremum of values will be the same both for the full space \mathcal{X} and for the restricted class $\mathcal{X} \setminus \mathcal{H}$. In other words, we now see that $S(k)$ can be obtained as supremum over the whole of \mathcal{X} , and the only issue, which may bother us a little, if the actual maximizers, if we find some, will belong to the singular hyperplane \mathcal{H} , or stay in $\mathcal{X} \setminus \mathcal{H}$. That we need to check at the end.

¹Of course, this allegedly holding continuity must be, and indeed will be shown easily later.

2.2. An estimation of the extremal value. Before proceeding let us stop for a little further analysis, establishing at least that $S(k)$ is bounded, because this will be needed in what follows.

Soundararajan [11] remarks that “the unfortunate inequality” $S(k) < 4/k$ holds. This is not completely obvious, but in fact the situation is even worse, namely, $S(k) < 4/(k + c \log k)$. This was essentially proved (without an explicit calculation of c) in the mentioned unpublished note of Soundararajan [10]. Together with the mentioned example $P(x) = x^{k+\ell}$, $\ell = c\sqrt{k}$, this shows that the value of $S(k)$ is between $4/(k + c' \log k)$ and $4/(k + c''\sqrt{k})$.

Proposition 1 (Soundararajan). *The extremal problem (1) is bounded by $4/k$. Moreover, we have $S(k) < \frac{4}{k + \log_2 k - 5}$ for all $k \geq 4$.*

Proof. Let us put $q(x) = Q(1 - x)$, and take $m > n/2$, $I(m) := \int_0^1 x^m q^2(x) dx$ and $J(n) := \int_0^1 x^n q'^2(x) dx$. Partial integration (using also $q(1) = Q(0) = 0$) and the Cauchy–Schwarz inequality yield

$$I(m) = \frac{-1}{m+1} \int_0^1 x^{m+1} 2q'(x)q(x) dx \leq \frac{2}{m+1} \sqrt{I(2m+2-n)J(n)}.$$

Starting from $m = k - 2$ and $n = k - 1$ and continuing by induction, we thus arrive at

$$\frac{I(m)}{J(n)} \leq \prod_{j=0}^N \left(\frac{2}{k + 2^j - 2} \right)^{2^{-j}} \cdot I(k + 2^{N+1} - 3)^{2^{-(N+1)}} J(n)^{-2^{-(N+1)}}.$$

Since $0 \leq I(\nu)$ is decreasing with ν , $I(k + 2^{N+1} - 3)$ converges with N and we can pass to the limit $N \rightarrow \infty$, then even take supremum with respect to q , and hence obtain

$$S(k) \leq (k-1) \prod_{j=0}^{\infty} \left(\frac{2}{k + 2^j - 2} \right)^{2^{-j}} = \frac{4}{\prod_{j=1}^{\infty} (k + 2^j - 2)^{2^{-j}}} = \frac{4}{k-2} \prod_{j=1}^{\infty} \left(1 + \frac{2^j}{k-2} \right)^{-2^{-j}}.$$

Observe that for $j > 1$ every single $k + 2^j - 2 \geq k + 1 > k$ in the denominator of the last but one expression, hence $S(k) < 4/k$ follows immediately.² We can even sharpen this estimate further. Let us denote the last product by $D := D(k)$ and define $\ell := \lfloor \log_2(k-2) \rfloor$. Then, by using $\log(1+x) > x - \frac{1}{2}x^2$ (for $x > 0$) we infer

$$\begin{aligned} \log D(k) &= - \sum_{j=1}^{\infty} \frac{\log \left(1 + \frac{2^j}{k-2} \right)}{2^j} < - \sum_{j=1}^{\ell} \frac{1}{2^j} \left(\frac{2^j}{k-2} - \frac{1}{2} \left(\frac{2^j}{k-2} \right)^2 \right) \\ &= - \frac{\ell}{k-2} + \frac{2^{\ell} - 1}{(k-2)^2} < - \frac{\ell - 1}{k-2}. \end{aligned}$$

Therefore, as $e^{-x} < \frac{1}{1+x}$ (for $x > 0$), we obtain

$$S(k) < \frac{4}{k-2} \cdot \exp \left(- \frac{\ell - 1}{k-2} \right) < \frac{4}{(k-2) \left(1 + \frac{\ell-1}{k-2} \right)} = \frac{4}{k + \ell - 3} \leq \frac{4}{k + \log_2 k - 5},$$

since $\ell \geq \log_2(k-2) - 1 \geq \log_2 k - 2$ for all $k \geq 4$. \square

²Note that we did not need anything more than $m, n \in \mathbb{N}$, $q(1) = 0$, and that the integrals $\int_0^1 x^{m-1} q^2(x) dx$ and $\int_0^1 x^n q'^2(x) dx$ exist finitely.

A further elementary observation is that for the Cauchy–Schwarz inequality to be precise, we should have $x^{m+1-n/2}q = cx^{n/2}q'$ in all the above applications of the Cauchy–Schwarz estimate (i.e., for all occurring values of m). This cannot hold for whatever choice of q for all m simultaneously. To have an about optimal estimate we may strive for having the Cauchy–Schwarz estimate sharp at the very first application, when $m = k - 2$ and $n = k - 1$, so $q = cq'$ follows, and then $q(x) = e^{cx}$. But even that is not a valid choice in our problem: $q(1) = 0$ prevents us taking q an exponential function as it can never be zero. In any case, the estimate of $S(k)$ above cannot be sharp.

2.3. A first analysis of the extremal problem. Now let us go a little further and not only implement the said change of normalization, but also write in $q(x) = Q(1 - x)$ as before in the estimations. Then the whole problem becomes

$$(5) \quad \max (k - 1) \left(\int_0^1 x^{k-2} q^2(x) dx \right) / \left(\int_0^1 x^{k-1} q'^2(x) dx \right) \quad \text{under condition } q(1) = 0,$$

understood in an appropriate function class \mathcal{X} , like, e.g., $C^1[0, 1]$.

Partial integration in the numerator and $q(1) = 0$ yields now the reformulation

$$(6) \quad S(k) = \sup_{q \in \mathcal{X}, q(1)=0, q \neq 0} \left(-2 \int_0^1 x^{k-1} q'(x) q(x) dx \right) / \left(\int_0^1 x^{k-1} q'^2(x) dx \right).$$

Certainly we want the denominator to be finite, so we assume that our function class is chosen so that for any $q \in \mathcal{X}$ this weighted square integral of q' converges. On the other hand, as mentioned several times before, this implies both the convergence of the numerator (and boundedness of the fraction by say $4/k$) and also that the positive, non-degenerate linear functional $I(q) := (k - 2)!P(1) = \int_0^1 q(t)t^{k-2}dt$ is well-defined, finite. So now we fix the largest function space we may deal with as

$$(7) \quad \mathcal{X} := \left\{ q : (0, 1] \rightarrow \mathbb{R} : q(x) = - \int_x^1 q'(t)dt, \int_0^1 x^{k-1} q'^2(x) dx < \infty \right\}.$$

The definition above means that q is an absolutely continuous function on each compact subinterval of $(0, 1]$, whence $q' \in L^1_{\text{loc}}(0, 1]$ and $q(x)$ exists as a Lebesgue integral of q' ; further q' is square-integrable with respect to the weight x^{k-1} .

3. EXISTENCE AND SMOOTHNESS OF MAXIMIZERS IN THE EXTREMAL PROBLEM

3.1. Existence of maximizing functions in the extremal problem. In this paper the role of k is fixed. Furthermore, it will be convenient for us to avoid repetitious use of $k - 2$ and $k - 1$, so throughout the rest of the paper except for the last section, Section 6, we will fix the notations for two further integer parameters. So we define

$$(8) \quad m := k - 2, \quad n := k - 1.$$

As it is explained above, we can discuss the optimization problem in the function space

$$(9) \quad \mathcal{Y} := \left\{ p(t) \in L^1_{\text{loc}}(0, 1] : \int_0^1 p^2(t)t^n dt < \infty \right\}, \quad \text{where } q(x) = - \int_x^1 p(t)dt.$$

Multiplying the occurring functions by $t^{n/2}$, we can even consider the space of functions $\varphi(t) := p(t)t^{n/2} = q'(t)t^{n/2}$, which then will be square-integrable on $[0, 1]$, so that $\varphi \in L^2[0, 1]$.

Next let us establish, how the functional to be maximized looks like over these spaces. On \mathcal{X} , on \mathcal{Y} and finally on $L^2[0, 1]$ we must consider the respective equivalent expressions

$$\begin{aligned}
& \frac{-2 \int_0^1 x^{k-1} q'(x) q(x) dx}{\int_0^1 x^{k-1} q'^2(x) dx} = \frac{2 \int_0^1 x^n p(x) \left(\int_x^1 p(t) dt \right) dx}{\int_0^1 p^2(x) x^n dx} \\
& = \frac{2 \int_0^1 \varphi(x) x^{n/2} \left(\int_x^1 \varphi(t) t^{-n/2} dt \right) dx}{\int_0^1 \varphi^2(x) dx} = \frac{2 \int_0^1 \int_0^1 \varphi(x) \varphi(t) \chi_{t>x}(x, t) x^{n/2} t^{-n/2} dt dx}{\int_0^1 \varphi^2(x) dx} \\
(10) \quad & = \frac{\int_0^1 \int_0^1 \varphi(x) \varphi(t) K(x, t) dt dx}{\int_0^1 \varphi^2(x) dx} \quad \text{with} \quad K(x, t) := \left(\frac{\min(x, t)}{\max(x, t)} \right)^{n/2},
\end{aligned}$$

the last step being a technical one to bring the kernel K to a symmetric form. So finally we find that

$$(11) \quad S(k) = \sup_{L^2[0,1] \setminus \{\mathbf{0}\}} \frac{\int_0^1 \int_0^1 \varphi(x) \varphi(t) K(x, t) dt dx}{\int_0^1 \varphi^2(x) dx} \quad \text{with} \quad K(x, t) := \left(\frac{\min(x, t)}{\max(x, t)} \right)^{n/2}.$$

Clearly, on $L^2[0, 1]$ the functional in (10) is defined everywhere except $\varphi = \mathbf{0}$ (the zero function), and is bounded by $4/k$, as proved before. Moreover, there is a clear homogeneity property: The ratio for any φ is equal to the ratio for any nonzero constant multiple $c\varphi$, hence the ratio is constant on all rays $\{c\varphi : c \in \mathbb{R}, c \neq 0\}$.

Therefore, the range of this quotient functional is clearly the same on the whole space $L^2[0, 1] \setminus \{\mathbf{0}\}$ as on $B \setminus \{\mathbf{0}\}$, B denoting the unit ball $B := \{\varphi \in L^2[0, 1] : \|\varphi\|_2 \leq 1\}$ (where the 2-norm of a function in $L^2[0, 1]$ is $\|\varphi\|_2 := (\int_0^1 \varphi^2(x) dx)^{1/2}$, as usual). Furthermore, actually already *on the unit sphere* $S := \{\varphi \in L^2[0, 1] : \|\varphi\|_2 = 1\}$ the functional must take on all the values of its range. However, on the unit sphere the denominator is exactly one, so now we can modify the formulation and write

$$S(k) = \sup_S \int_0^1 \int_0^1 \varphi(x) \varphi(t) K(x, t) dt dx = \sup_B \int_0^1 \int_0^1 \varphi(x) \varphi(t) K(x, t) dt dx.$$

Moreover, it is clear that in this last formulation $S(k)$ is taken by a maximizer function $\varphi \in L^2[0, 1]$ iff there is a maximum at some $\varphi \in S$ iff there is a maximum on B (in which case again any maximum must belong to S). So any maximizer φ in the original formulation is maximizer together with all the ray $\{c\varphi\}$ of its homothetic copies, and in the new formulation this maximizer occurs exactly with $c = \pm 1/\|\varphi\|_2$, i.e., at the unit norm elements of the given ray.

This reformulation furnishes us the access to settle the existence question of some maximizer. In our formulation of the extremal problem all functions are real-valued, only for the next two propositions (spectral theory), and for the sake of being precise, we shall need complex-valued functions.

Proposition 2. *Let*

$$K(x, y) := \left(\frac{\min(x, y)}{\max(x, y)} \right)^{n/2},$$

and define the Fredholm-type operator

$$(12) \quad T : L^2([0, 1]; \mathbb{C}) \rightarrow L^2([0, 1]; \mathbb{C}) \quad (T\varphi)(x) := \int_0^1 \varphi(y) K(x, y) dy.$$

Then T is a compact, positive, self-adjoint operator on the complex Hilbert space $L^2([0, 1]; \mathbb{C})$, maps real-valued functions into real-valued ones, and preserves positivity.

Proof. Since $K \in L^\infty([0, 1] \times [0, 1])$, T is compact, see [9, §97]. Since $0 \leq K \leq 1$ and K is symmetric, the other two properties follow evidently. \square

Proposition 3. $S(k)$ is attained as a maximum by some maximizing function $\varphi \in L^2[0, 1]$.

Equivalently,

$$S(k) = \frac{-2 \int_0^1 x^{k-1} q'(x) q(x) dx}{\int_0^1 x^{k-1} q'^2(x) dx}$$

for some appropriate $q \in \mathcal{X}$ with $q(1) = 0$ and $q \not\equiv 0$.

Proof. Consider the operator T as in (12).

$$(13) \quad A(\varphi, \psi) := \int_0^1 \int_0^1 \varphi(y) \overline{\psi}(x) K(x, y) dx dy = \langle T\varphi, \psi \rangle,$$

which is a sesquilinear form on $L^2([0, 1]; \mathbb{C})$. By [9, §93] we have

$$\|T\| := \sup_{\|\varphi\|_2 \leq 1} \|T\varphi\| = \sup_{\|\varphi\|_2 \leq 1} |\langle T\varphi, \varphi \rangle| = \sup_{\|\varphi\|_2 \leq 1} A(\varphi, \varphi).$$

Since T is compact, positive and self-adjoint, all of its eigenvalues are nonnegative, and the eigenvalues can be ordered in a decreasing null-sequence (λ_j) , $\lambda_1 > \dots > \lambda_j > \dots$, $\lambda_j \rightarrow 0$ ($j \rightarrow \infty$), and we also have

$$\|T\| = \max\{\lambda : \lambda \text{ is an eigenvalue of } T\} =: \lambda_1.$$

Since T leaves the subspace of real-valued functions invariant, for any eigenvalue $\lambda \in \mathbb{R}$ of T there is a real-valued eigenfunction³. Summing up, $\|T\| = \lambda_1$, and there exists some (nonzero) eigenfunction $\varphi \in L^2[0, 1]$ satisfying $\|\varphi\| = 1$ and $\lambda_1 = \|T\| = A(\varphi, \varphi)$, yielding a maximizer for $A(\varphi, \varphi)$ as asserted. \square

Remark 4. The above proof yields also the following important information: $S(k)$ is the largest eigenvalue λ_1 of T , and any (normalized) eigenfunction φ of T belonging to λ_1 is a maximizer; moreover, the only maximizers are nonzero eigenfunctions of T corresponding to $\lambda_1 = \|T\|$.

Indeed, as T is compact and self-adjoint, there is an orthonormal basis (e_j) in $L^2[0, 1]$ that consists of eigenfunctions of T . Let $\mathbf{0} \neq \varphi \in L^2[0, 1]$ be not an eigenfunction to the eigenvalue λ_1 . Then $\varphi = \sum_{j=1}^{\infty} \langle \varphi, e_j \rangle e_j$ and

$$\langle T\varphi, \varphi \rangle = \sum_{j=1}^{\infty} \lambda_j |\langle \varphi, e_j \rangle|^2 < \lambda_1 \|\varphi\|_2^2$$

by Parseval's identity, where for the strict inequality " $<$ " we have used that for some $j > 1$ we have $|\langle \varphi, e_j \rangle| > 0$, while $\lambda_j < \lambda_1$.

Next we turn to smoothness properties of eigenfunctions of T .

³By the way, since K is strictly positive, so is the operator T , hence one knows from Perron–Frobenius theory (see [7, Sec. 4.2]) that $\ker(T - \lambda_1)$ is one-dimensional, but this will be proved later anyway, as we determine this subspace exactly.

3.2. Smoothness of maximizers and maximizers in $C[0, 1]$. The above formulation also provides us a direct access to further smoothness statements.

Lemma 5. *The Fredholm-type operator T defined in (12) maps $L^2[0, 1]$ to the subspace $C_0(0, 1]$ of continuous functions with value 0 at 0.*

Proof. Since $L^2[0, 1] \subset L^1[0, 1]$, and $0 \leq K(x, y) \leq 1$, the expression $(T\varphi)(x) = \int_0^1 \varphi(y)K(x, y)dy$ is an integral with a uniform majorant $|\varphi(y)| \in L^1[0, 1]$ of the integrands. Hence by the Lebesgue Dominated Convergence Theorem, it suffices to take the pointwise limit under the integral sign. When $x \rightarrow x_0$, this gives for all $y > 0$ $\lim_{x \rightarrow x_0} K(x, y) = K(x_0, y)$, while for $y = 0$ we have $K(x, 0) = K(x_0, 0) = 0$ identically. (Essentially, we have used only separate continuity of K on $[0, 1] \times [0, 1]$.) Thus $\lim_{x \rightarrow x_0} (T\varphi)(x) = \int_0^1 \varphi(y) \lim_{x \rightarrow x_0} K(x, y)dy = \int_0^1 \varphi(y)K(x_0, y)dy = (T\varphi)(x_0)$, i.e., $T\varphi \in C[0, 1]$. By definition, $K(0, y) = 0$ for all $y \in [0, 1]$, hence for every $\varphi \in L^2[0, 1]$ we have $(T\varphi)(0) = 0$. \square

Corollary 6. *All eigenfunctions φ of the Fredholm-type operator T defined in (12) are continuous and fulfill $\varphi(0) = 0$.*

Equivalently, in the function space \mathcal{X} defined in (7) all the functions $q(x) = -\int_x^1 q'(t)dt = -\int_x^1 t^{-n/2}\varphi(t)dt$ corresponding to eigenfunctions φ of T satisfy $\varphi(x) = x^{n/2}q'(x) \in C_0(0, 1]$ and thus $x^{n/2-1}q(x) \in C_0(0, 1]$.

Proof. All eigenfunctions lie in the range of the operator T , hence belong to $C_0(0, 1]$ in view of Lemma 5.

Recall that the correspondence between $L^2[0, 1]$ and our spaces \mathcal{Y} and \mathcal{X} was given by $\varphi(x) = x^{n/2}p(x) = x^{n/2}q'(x)$. Thus for $\varphi \in L^2[0, 1]$, an eigenfunction of T , we obtain for the corresponding q that $x^{n/2}q'(x) \in C_0(0, 1]$, whence also $q' \in C(0, 1]$ follows. Moreover, $\lim_{x \rightarrow 0+} x^{n/2}q'(x) = \lim_{x \rightarrow 0+} \varphi(x) = \varphi(0) = 0$, providing a continuous extension of $x^{n/2}q'(x)$ even to 0. Now writing $q(x) = -\int_x^1 q'(t)dt$ yields $q \in C(0, 1]$. While for $x \rightarrow 0+$ we obtain that

$$\lim_{x \rightarrow 0+} x^{n/2-1}q(x) = \lim_{x \rightarrow 0+} x^{n/2-1} \left(-\int_x^1 o(1)t^{-n/2}dt \right) = \lim_{x \rightarrow 0+} o(1) \frac{2}{n-2} = 0,$$

hence $x^{n/2-1}q(x) \in C_0(0, 1]$. \square

Although K is not everywhere continuous $[0, 1] \times [0, 1]$ the operator T can be approximated by compact operators given by continuous kernels.

Proposition 7. *The operator T restricted to $C[0, 1]$ is a compact $C[0, 1] \rightarrow C[0, 1]$ operator with exactly the same eigenvalues and eigenfunctions as on $L^2[0, 1]$.*

Proof. Let $f_j : [0, 1] \times [0, 1] \rightarrow [0, 1]$ be continuous with $f_j(x, y) = 0$ if $x, y \leq \frac{1}{2j}$ and $f_j(x, y) = 1$ if $\max(x, y) \geq \frac{1}{j}$. Then $f_j K$ is continuous, hence the integral operator T_j with kernel $f_j K$ is compact, see, e.g., [9, §90]. It is easy to see that $T_j \rightarrow T$ in the operator norm (over $C[0, 1]$), hence T itself is compact, see [9, §76].

By Proposition 6 all eigenfunctions of T on $L^2[0, 1]$ belong also to $C_0(0, 1]$, hence remain eigenfunctions when T is considered as $C[0, 1] \rightarrow C[0, 1]$. The converse is obvious: Every continuous eigenfunction is of course also an eigenfunction from $L^2[0, 1]$. In particular, the set of eigenvalues are also exactly the same when considered in these two spaces. \square

Remark 8. *One can show that the norm of T as an operator on $C[0, 1]$ is*

$$(14) \quad \|T\|_{C[0,1]} = \left(\frac{4}{n+2}\right)^{\frac{n}{n-2}} = \left(\frac{4}{k+1}\right)^{\frac{k-1}{k-3}}.$$

In fact, since T is positivity preserving, we have $\|T\|_{C[0,1]} = \|T\mathbf{1}\|_{\infty}$, where $\mathbf{1}$ is the constant 1 function. Easy calculation shows that $(T\mathbf{1})(x) = \frac{4nx}{n^2-4} - \frac{nx^{n/2-1}}{n-2}$ and that this function has maximum at $x = (\frac{2}{n+2})^{2/(n-2)}$. Since we already know $\|T\|_{L^2[0,1]} = \lambda_1$, and in general $\lambda_1 \leq \|T\|_{C[0,1]}$, we obtain that the maximum of $A(\varphi, \varphi)$ with $\|\varphi\|_2 = 1$, i.e., $S(k)$ is smaller than the constant in (14) above.

3.3. Differentiability of maximizers. We now push further the smoothness statements from the last subsection. We need some preparations, and define the following auxiliary functions

$$(15) \quad \kappa(x, y) := \begin{cases} \frac{\min(x,y)}{\max(x,y)} & \text{if } (0,0) \neq (x,y) \in [0,1]^2, \\ 0 & \text{if } (x,y) = (0,0), \end{cases}$$

and

$$(16) \quad \omega(a, b) := \begin{cases} \frac{\sum_{j=0}^{n-1} a^{j/2} b^{(n-1-j)/2}}{\sqrt{b} + \sqrt{a}} & \text{if } (0,0) \neq (a,b) \in [0,1]^2, \\ 0 & \text{if } (a,b) = (0,0). \end{cases}$$

With these notations we have for every $0 < x, y \leq 1$ the formula

$$(17) \quad \frac{n}{2} K(x, y) = \omega(\kappa(x, y), \kappa(x, y)) \kappa(x, y),$$

which also holds for x or y being 0, with both sides vanishing.

Now note that $0 \leq \omega(a, b) \leq n \max(a, b)^{n/2-1} \leq n$. Furthermore, observe that for $0 \leq a, b \leq 1$ (even if both are zero) we have $b^{n/2} - a^{n/2} = (\sqrt{b} - \sqrt{a}) \sum_{j=0}^{n-1} a^{j/2} b^{(n-1-j)/2} = (b-a)\omega(a, b)$. Hence for any $y, x, x' \in [0, 1]$ we can write

$$(18) \quad K(x', y) - K(x, y) = \omega(\kappa(x', y), \kappa(x, y)) \cdot (\kappa(x', y) - \kappa(x, y)).$$

Fix $x > 0$. Denoting $\Delta := x' - x > 0$, we also have

$$(19) \quad \kappa(x', y) - \kappa(x, y) = \begin{cases} \frac{y}{x'} - \frac{y}{x} = -\frac{y\Delta}{xx'} & \text{if } y < x < x', \\ \frac{y}{x'} - \frac{y}{x} = \frac{y^2 - xx'}{yx'} & \text{if } x \leq y \leq x', \\ \frac{x'}{y} - \frac{x}{y} = \frac{\Delta}{y} & \text{if } x < x' < y, \end{cases}$$

whence

$$(20) \quad |\kappa(x', y) - \kappa(x, y)| = \begin{cases} \frac{y}{x} - \frac{y}{x'} = \frac{y\Delta}{xx'} \leq \frac{\Delta}{x'} \leq \frac{\Delta}{x} & \text{if } y < x < x', \\ \left| \frac{x}{y} - \frac{x'}{y} \right| = \frac{|xx' - y^2|}{yx'} \leq \frac{\Delta}{y} \leq \frac{\Delta}{x} & \text{if } x \leq y \leq x', \\ \frac{x'}{y} - \frac{x}{y} = \frac{\Delta}{y} \leq \frac{\Delta}{x'} \leq \frac{\Delta}{x} & \text{if } x < x' < y. \end{cases}$$

Lemma 9. *The operator T maps $L^2[0, 1]$ to the space $C^1(0, 1]$. Furthermore, for $\varphi \in L^2[0, 1]$ and $x \in (0, 1]$*

$$(21) \quad (T\varphi)'(x) = \int_0^1 \varphi(y) \frac{\partial}{\partial x} K(x, y) dy = -\frac{n}{2x} (T\varphi)(x) + nx^{n/2-1} \int_x^1 \frac{\varphi(y)}{y^{n/2}} dy.$$

2(Sz.R.): C
tion in the n
of the second

Proof. The two expressions given for $(T\varphi)'(x)$ in (21) are easily seen to be equal, so the proof hinges upon showing that $(T\varphi)'(x)$ equals any one of them.

For any $0 \leq x < x' \leq 1$ using (18) we can write

$$(22) \quad \frac{(T\varphi)(x') - (T\varphi)(x)}{x' - x} = \int_0^1 \varphi(y) \omega(\kappa(x', y), \kappa(x, y)) \frac{\kappa(x', y) - \kappa(x, y)}{x' - x} dy.$$

We fix $x_0 > 0$ and take either $x = x_0$ and $x' \rightarrow x_0+$, or $x' = x_0$ and $x \rightarrow x_0-$. In any case, by (20) and $0 \leq \omega \leq n$ we have the Lebesgue integrable majorant $n|\varphi(y)|/x_0$ of the integrand, thus limit and integral can be interchanged.

For example in the case $x' \rightarrow x_0+$ taking into account (17) we are led to

$$(23) \quad \begin{aligned} \lim_{x' \rightarrow x_0} \frac{(T\varphi)(x') - (T\varphi)(x_0)}{x' - x_0} &= \int_0^1 \varphi(y) \omega(\kappa(x_0, y), \kappa(x_0, y)) \frac{\partial}{\partial x} \kappa(x, y) \Big|_{x=x_0} dy \\ &= \int_0^1 \varphi(y) \omega(\kappa(x_0, y), \kappa(x_0, y)) \operatorname{sgn}(y - x_0) \frac{\kappa(x_0, y)}{x_0} dy \\ &= \frac{n}{2x_0} \int_0^1 \varphi(y) K(x_0, y) \operatorname{sgn}(y - x_0) dy = \frac{n}{2x_0} \left(\int_{x_0}^1 \varphi(y) K(x_0, y) dy - \int_0^{x_0} \varphi(y) K(x_0, y) dy \right) \\ &= \frac{n}{x_0} \left(\int_{x_0}^1 \varphi(y) K(x_0, y) dy - \frac{1}{2} (T\varphi)(x_0) \right), \end{aligned}$$

where we have used $\frac{\partial}{\partial x} \kappa(x, y) = \frac{\partial}{\partial x} (x/y) = 1/y$ for $y > x$ and $\frac{\partial}{\partial x} \kappa(x, y) = \frac{\partial}{\partial x} (y/x) = -y/x^2$ for $x < y$. When substituting the definition of K in the above, we obtain all the asserted formulas. Note that in case $y = x_0$, one sided derivatives of $\kappa(x_0, y)$ still exist (and are equal to the limits from the respective side) but the existence and value of the limit at one exceptional point does not interfere the value of the integral, therefore we have just put 0 for the value of $\frac{\partial}{\partial x} \kappa(x, x_0) \Big|_{x=x_0}$ here.

When $x \rightarrow x_0- = x'-$, the calculation is entirely the same.

The integrals on the right hand side of (21) are of course integrals of integrable functions, and as such, are continuous in function of the limits of integration. Therefore, continuity of $(T\varphi)'$ on $(0, 1]$ also follows. \square

Remark 10. When $x_0 = 0$, only the right hand side derivative can be considered and thus we take $x_0 = x = 0$ and $x' \rightarrow 0+$. Also, $(T\varphi)(0) = 0$ and $K(0, y) = 0$, hence the consideration of the differential reduces to

$$(24) \quad \lim_{x' \rightarrow 0+} \frac{(T\varphi)(x')}{x'} = \lim_{x' \rightarrow 0+} \frac{1}{x'} \int_0^1 \varphi(y) K(x', y) dy = \lim_{x' \rightarrow 0+} \int_0^1 \varphi(y) \frac{\kappa^{n/2}(x', y)}{x'} dy,$$

which, however, cannot be handled for general $\varphi \in L^2[0, 1]$ or not even for $\varphi \in C_0(0, 1]$, and can be well estimated only if we use something more on φ . See Corollary 12 below.

Proposition 11. The operator T maps $L^2[0, 1]$ to the space of absolutely continuous functions with bounded total variation. Moreover, for the total variation of $T\varphi$ we have $V(T\varphi, [0, 1]) \leq 2\|\varphi\|_1$.

Proof. We already know that $T\varphi \in C^1(0, 1]$, so the total variation, whether finite or infinite, can be computed as $V(T\varphi, [0, 1]) = \int_0^1 |(T\varphi)'|$. Now the first formula from (21)

furnishes

$$\begin{aligned} V(T\varphi, [0, 1]) &= \int_0^1 |(T\varphi)'(x)| dx = \int_0^1 \left| \int_0^1 \varphi(y) \frac{\partial}{\partial x} K(x, y) dy \right| dx \\ &\leq \int_0^1 \int_0^1 |\varphi(y)| \left| \frac{\partial}{\partial x} K(x, y) \right| dy dx = \int_0^1 |\varphi(y)| \left(\int_0^1 \left| \frac{\partial}{\partial x} K(x, y) \right| dx \right) dy = 2\|\varphi\|_1, \end{aligned}$$

as for all $0 < y < 1$ fixed we have $\int_0^1 \left| \frac{\partial}{\partial x} K(x, y) \right| dx = V(K(\cdot, y), [0, 1]) = 2$. \square

Corollary 12. *If ψ lies in the range of T , then $T\psi \in C^1[0, 1]$, and $(T\psi)'(0) = 0$. In particular if φ is an eigenfunction of T , then $\varphi \in C^1[0, 1]$, and $\varphi'(0) = 0$.*

Proof. We have to calculate the limit in (24) for $\psi := T\varphi$ in place of φ . Recall from the above that then $\psi = T\varphi \in C_0(0, 1]$, in particular $\psi(0) = (T\varphi)(0) = 0$, and $\psi \in C^1(0, 1]$, $V(\psi, [0, 1]) \leq 2\|\varphi\|_1$. The second mean value theorem and integration by parts yield with some appropriate $z := z_{x'} \in (0, x')$

$$\begin{aligned} \frac{1}{x'} \int_0^1 \psi(y) K(x', y) dy &= \frac{1}{x'} \left\{ \int_0^{x'} \psi(y) K(x', y) dy + \int_{x'}^1 \psi(y) \frac{x'^{n/2}}{y^{n/2}} dy \right\} \\ &= \psi(z_{x'}) K(x', z_{x'}) + \left\{ \left[\psi(y) \frac{-x'^{n/2-1}}{n/2-1} y^{1-n/2} \right]_{x'}^1 + \int_{x'}^1 \psi'(y) \frac{x'^{n/2-1}}{n/2-1} y^{1-n/2} dy \right\}, \end{aligned}$$

so the first term tends to $\psi(0) = 0$ when $x' \rightarrow 0+$. The term in the square bracket contributes $x'^{n/2-1} \frac{-\psi(1)}{n/2-1} + \frac{\psi(x')}{n/2-1}$, and as $x' \rightarrow 0+$ and $\psi(x') \rightarrow \psi(0) = 0$, both terms converge to 0. Finally, for the integral Proposition 11 gives that $\psi' \in L^1[0, 1]$, while the product of the further factors stays bounded uniformly for all $x', y \in [0, 1]$, as the integral runs only through values $y \geq x'$. That is, we can again use the Lebesgue Dominated Convergence Theorem and calculate the limit by moving it under the integral sign. Furthermore, the pointwise limit of the expression is zero for all fixed y , whence the assertion follows. \square

4. SOLVING THE MAXIMIZATION PROBLEM

4.1. Setting up a differential equation for potential extremal functions. By the previous section we know that our maximization problem has a solution, and we also saw that maximizers are sufficiently smooth. We can now set up a differential equation to find maximizers, or which is essentially equivalent, to find the eigenfunctions of T .

Proposition 13. *Let $\varphi \in L^2[0, 1]$ be an eigenfunction of T corresponding to the eigenvalue $\lambda > 0$. Then φ is continuous on $[0, 1]$, infinitely often differentiable on $(0, 1]$. The function $q(x) = -\int_x^1 \varphi(y) y^{-n/2} dy$ satisfies $q(1) = 0$ and the differential equation*

$$(25) \quad q''(x) + \frac{n}{x} q'(x) + \frac{b}{x} q(x) = 0 \quad \left(x \in (0, 1] \right), \quad \text{where } b := \frac{n}{\lambda} > 0.$$

Conversely, let $\lambda > 0$ and suppose that q is a nonzero, $C^2(0, 1]$ solution of the differential equation above with $q(1) = 0$. If $\varphi(x) = x^{n/2} q'(x)$ extends continuously to 0 with $\lim_{x \rightarrow 0+} x^{n/2} q'(x) = 0$, then φ is an eigenfunction of T corresponding to the eigenvalue $\lambda > 0$.

Proof. If $\varphi \in L^2[0, 1]$ is an eigenfunction of T for the eigenvalue $\lambda > 0$, then it belongs to the range of T , hence is continuous and continuously differentiable on $(0, 1]$ by Lemma 9.

Substituting $T\varphi = \lambda\varphi$ in (21) we obtain

$$\lambda\varphi'(x) = -\frac{n}{2x}\lambda\varphi(x) + nx^{n/2-1} \int_x^1 \frac{\varphi(y)}{y^{n/2}} dy.$$

As the right-hand side is differentiable, we can differentiate also the left-hand side showing $\varphi \in C^2(0, 1]$. We substitute $x^{-n/2}\varphi(x) = q'(x)$ and $\varphi(x) = x^{n/2}q'(x)$ and obtain

$$\frac{d}{dx} (\lambda x^{n/2} q'(x)) = -\frac{n}{2x} \lambda x^{n/2} q'(x) + nx^{n/2-1} \int_x^1 q'(y) dy,$$

and hence

$$\frac{d}{dx} (\lambda x^{n/2} q'(x)) = -\lambda \frac{n}{2} x^{n/2-1} q'(x) - nx^{n/2-1} q(x).$$

Differentiation yields

$$\lambda x^{n/2} q''(x) + \lambda \frac{n}{2} x^{n/2-1} q'(x) = -\frac{\lambda n}{2x} x^{n/2} q'(x) - nx^{n/2-1} q(x),$$

and then by rearranging we obtain

$$\lambda x^{n/2} q''(x) + \lambda \frac{n}{x} x^{n/2} q'(x) + nx^{n/2-1} q(x) = 0.$$

Division by $\lambda x^{n/2}$ thus leads to the asserted differential equation (25).

To see the converse we set $\psi = T\varphi$. Note that then $\psi \in C_0(0, 1]$ according to Lemma 5. Then $\varphi(x) = x^{n/2}q'(x)$ entails

$$q''(x) = -\frac{n}{2}x^{-n/2-1}\varphi(x) + x^{-n/2}\varphi'(x),$$

so that using the assumption that q solves (25) we obtain

$$-\frac{n}{2}x^{-n/2-1}\varphi(x) + x^{-n/2}\varphi'(x) + \frac{n}{x}x^{-n/2}\varphi(x) + \frac{n/\lambda}{x} \left(- \int_x^1 y^{-n/2}\varphi(y) dy \right) = 0,$$

and thus also

$$\frac{n}{2}\lambda x^{-n/2-1}\varphi(x) + \lambda x^{-n/2}\varphi'(x) - \frac{n}{x} \int_x^1 y^{-n/2}\varphi(y) dy = 0.$$

By (21) with $\psi = T\varphi$ we also have

$$x^{-n/2}\psi'(x) = -\frac{n}{2}x^{-n/2-1}\psi(x) + \frac{n}{x} \int_x^1 \frac{\varphi(y)}{y^{n/2}} dy,$$

so

$$\frac{n}{2}\lambda x^{-n/2-1}\varphi(x) + \lambda x^{-n/2}\varphi'(x) - x^{-n/2}\psi'(x) - \frac{n}{2}x^{-n/2-1}\psi(x) = 0.$$

If we multiply by x^n , we obtain for all $x > 0$

$$0 = \frac{n}{2}\lambda x^{n/2-1}\varphi(x) + \lambda x^{n/2}\varphi'(x) - x^{n/2}\psi'(x) - \frac{n}{2}x^{n/2-1}\psi(x) = \frac{d}{dx} (x^{n/2}\lambda\varphi(x) - x^{n/2}\psi(x)).$$

Since $\varphi, \psi \in C[0, 1]$, $x^{n/2}(\lambda\varphi(x) - \psi(x))$ must vanish at 0, whence $\lambda\varphi = \psi$ follows. \square

Thus the solution of the maximization problem is reduced to solving the homogeneous second order ordinary differential equation (25), and to finding the feasible values of λ . Next we solve this equation and analyze some properties of the solutions.

4.2. Bessel functions and Bessel's differential equation. Recall Bessel's differential equation

$$(26) \quad y''(x) + \frac{1}{x}y'(x) + \left(1 - \frac{\nu^2}{x^2}\right)y(x) = 0,$$

for some fixed parameter $\nu \in \mathbb{R}$. The Bessel function J_ν (of the first kind) is a solution of this equation, [2, (6.71), p. 115]. Notice that $J_{-\nu}$ is also a solution, but for ν integer J_ν and $J_{-\nu}$ are linearly dependent, in fact $J_{-\nu} = (-1)^\nu J_\nu$. So for $\nu \in \mathbb{N}$ another, linearly independent solution is needed, which is provided by the Bessel functions Y_ν (of the second kind), obtained as

$$Y_\nu(x) := J_\nu(x) \int \frac{dx}{xJ_\nu^2(x)},$$

where specifying the limits of integration is equivalent to fix some primitive of the integrand, and the integration limit cannot be at 0 where $Y_\nu(x)$ is divergent in the order $x^{-\nu}$, see [2, (6.73), (6.74)]. Then for $\nu \in \mathbb{N}$ the general solution of equation (26) is a linear combination $c_1J_\nu + c_2Y_\nu$, see [2, §102].

4.3. Solution of the differential equation. Bowman computes, see [2, (6.80), p. 117], what happens if we consider the transformed, substituted functions $u(x) := x^\alpha y(\beta x^\gamma)$, where y satisfies the Bessel equation (26), and establishes that then the new functions $u(x)$ will be the general solutions of the transformed equation

$$(27) \quad u''(x) - \frac{2\alpha - 1}{x}u'(x) + \left(\beta^2\gamma^2x^{2\gamma-2} + \frac{\alpha^2 - \nu^2\gamma^2}{x^2}\right)u(x) = 0.$$

If we choose here the parameters $\nu := m$, $\alpha := -m/2 = 1 - k/2$, $\beta := 2\sqrt{n/\lambda}$ and $\gamma := 1/2$ (where $n := k - 1 = m + 1$ and $m := k - 2$ as fixed above in (8)), then the equation (27) becomes exactly (25). Thus we obtain that for any fixed values of $m := k - 2$ and $\lambda > 0$, there is a one-to-one correspondence between the solutions q of (25) and y of (26) given by $q(x) := x^\alpha y(\beta x^\gamma) = x^{1-k/2}y(2\sqrt{n/\lambda}\sqrt{x})$.

Corollary 14. *Every solution q of (25) is a linear combination of transformed Bessel functions from the above, i.e.,*

$$q(x) = c_1x^{-m/2}J_m(2\sqrt{n/\lambda}\sqrt{x}) + c_2x^{-m/2}Y_m(2\sqrt{n/\lambda}\sqrt{x}), \quad (m = n - 1 = k - 2).$$

4.4. Some analysis of the occurring Bessel type functions. Before proceeding, let us note one important thing. Not all solutions of the differential equation are relevant for us, because the resulting q' must have finite weighted square integral, i.e., $\varphi(x) = x^{n/2}q'(x)$ has to belong to $L^2[0, 1]$ (and even be continuous on $[0, 1]$, according to Proposition 13).

According to the second formula of [1, 9.1.30] (with the choice $k = 1$ and $\nu = m$ there) we have

$$\begin{aligned} & \left(\frac{1}{x} \frac{d}{dx}\right) (x^{-m}c_1J_m(x) + c_2Y_m(x)) \\ &= -x^{-m-1} (c_1J_{m+1}(x) + c_2Y_{m+1}(x)) = -x^{-n} (c_1J_n(x) + c_2Y_n(x)), \end{aligned}$$

therefore we obtain for a solution q of (25) that

$$\begin{aligned} q'(x) &= \frac{d}{dx} \left(c_1x^{-m/2}J_m(2\sqrt{n/\lambda}\sqrt{x}) + c_2x^{-m/2}Y_m(2\sqrt{n/\lambda}\sqrt{x}) \right) \\ &= -\frac{1}{2\sqrt{n/\lambda}}x^{1/2}x^{-n/2} \left(c_1J_n(2\sqrt{n/\lambda}\sqrt{x}) + c_2Y_n(2\sqrt{n/\lambda}\sqrt{x}) \right). \end{aligned}$$

So that

$$(28) \quad \varphi(x) = x^{n/2}q'(x) = -\frac{\sqrt{x}}{2\sqrt{n/\lambda}} \left(c_1 J_n(2\sqrt{n/\lambda}\sqrt{x}) + c_2 Y_n(2\sqrt{n/\lambda}\sqrt{x}) \right).$$

To see when such a φ may actually be an eigenfunction of T , we first find out when it belongs to $C_0(0, 1]$. First, see [2, (1.2)]

$$(29) \quad J_\nu(x) = x^\nu \sum_{j=0}^{\infty} \frac{(-1)^j}{2^{\nu+2j}} \frac{x^{2j}}{j!(\nu+j)!}, \quad \text{whence also } J_\nu(x) \sim \frac{1}{2^\nu \nu!} x^\nu \quad (x \rightarrow 0+),$$

so in particular J_n is continuous on $[0, \infty)$, and for any $c_1 \in \mathbb{R}$ the part $\frac{\sqrt{x}}{2\sqrt{n/\lambda}} c_1 J_n(2\sqrt{n/\lambda}\sqrt{x})$ belongs to $C_0(0, 1] \subset L^2[0, 1]$. Second, $Y_n(x) \asymp x^{-n}$ ($x \rightarrow 0+$) [2, p. 116], entailing that for $c_2 \neq 0$ $\frac{\sqrt{x}}{2\sqrt{n/\lambda}} c_2 Y_n(2\sqrt{n/\lambda}\sqrt{x}) \asymp x^{-(n-1)/2}$, and this function is not even bounded near 0, if $n > 1$ (i.e., when $k = n - 1 > 2$). If $n = 1$, i.e., $k = 2$, this function is not vanishing at 0, a condition that is necessary for an eigenfunction of T by Lemma 5. So from Corollary 6 we obtain that φ in (28) belongs to $C_0(0, 1]$ (and thus may be a candidate for being an eigenfunction of T) if and only if $c_2 = 0$.

As a consequence of this and of Proposition 13 we obtain the following.

Corollary 15. *Consider the operator T from (12), and let $\lambda > 0$. Then $\lambda > 0$ is an eigenvalue of T if and only if*

$$J_m \left(2\sqrt{n/\lambda} \right) = 0.$$

In this case

$$\varphi(x) = x^{n/2}q'(x) = c_1 x^{1/2} J_n(2\sqrt{n/\lambda}\sqrt{x}), \quad c_1 \neq 0$$

are the only eigenfunctions corresponding to λ .

For given m let us denote the roots of J_m by $\alpha_{m,r}$ ($r \in \mathbb{N}$) ordered increasingly. At this stage it is in order to recall the following about the zeros of Bessel functions. We have $\alpha_{m,r} \rightarrow \infty$ ($r \rightarrow \infty$), and for rather large values the roots $\alpha_{m,r}$ of J_m are very well distributed, as essentially there falls one root in each interval of length π . However, for fixed m the increasing sequence of zeros $(\alpha_{m,r})$ starts only with $\alpha_{m,1} \sim m + cm^{1/3}$, with $c = 1.8557571\dots$, see [1, 9.5.14, p. 341]. Let us introduce the notation

$$\lambda_{m,r} := \lambda_r := 4(m+1)/\alpha_{m,r}^2$$

and

$$q_r(x) := q_{m,r}(x) := x^{-m/2} J_m \left(2\sqrt{(m+1)/\lambda_{m,r}} \cdot \sqrt{x} \right) = x^{-m/2} J_m \left(\alpha_{m,r} \sqrt{x} \right).$$

By putting everything together we obtain the following result.

Theorem 16. *For the extremal problem (1) we have*

$$(30) \quad S(k) = \lambda_1 = \frac{4(k-1)}{\alpha_{k-2,1}^2} = \frac{4}{k + 2ck^{1/3} + O(1)}$$

with $\alpha_{k-2,1}$ the first root of the order $k-2$ Bessel function $J_{k-2}(x)$, and the constant $c = 1.8557571\dots$

The only extremal functions for the formulation (5) are nonzero constant multiples of

$$q_1(x) = x^{-(k-2)/2} J_{k-2}(\alpha_{k-2,1}\sqrt{x}).$$

Proof. Uniqueness follows from Remark 4 and Corollary 15. For q_1 being a maximizer for (5) it remains only to show that

$$\int_0^1 q_1(y)y^n dy \neq 0.$$

But this follows since q_1 is (strictly) positive all over $(0, 1)$, $\alpha_{k-2,1}$ being the very first zero of the Bessel function J_{k-2} . \square

5. POWER SERIES

The above settles the issue of the best weight P —and also the order of $S(k)$ —in the GPY method. However, not all weight functions are easy to handle, and a Bessel function—even if an analytic function with relatively strongly convergent series expansion—may be unmanageable, at least in our current technical abilities. We will discuss, using the classical series expansion of J_m , how it may work in this context. Actually, not too well.

With the notations from the above for q_r , $\alpha_{m,r}$ etc., (29) yields

$$q_r(x) = x^{-m/2}\beta^{-m}J_m(2\beta\sqrt{x}) = \sum_{j=0}^{\infty} \frac{(-b)^j}{j!(m+j)!}x^j, \quad \text{with } \beta := \sqrt{b}, \quad b = \beta^2 = \frac{\alpha_{m,r}^2}{4},$$

Then we can evaluate the functionals

$$G(q_r) = \int_0^1 x^{m+1}q_r'^2(x)dx, \quad F(q_r) = \int_0^1 (m+1)x^m q_r^2(x)dx = (-2) \int_0^1 x^{m+1}q_r(x)q_r'(x)dx.$$

Computations with the power series provide

$$G(q_r) = b^2 \sum_{j,\ell=0}^{\infty} \frac{(-b)^{j+\ell}}{(m+j+\ell+2)j!\ell!(m+j+1)(m+\ell+1)!},$$

and

$$F(q_r) = 2b \sum_{j,\ell=0}^{\infty} \frac{(-b)^{j+\ell}}{(m+j+\ell+2)j!\ell!(m+j)(m+\ell+1)!}.$$

With some reformulations we can also write

$$\begin{aligned} \frac{F(q_r)}{G(q_r)} &= \frac{2}{b} \frac{\sum_{\nu=0}^{\infty} \frac{(-b)^\nu}{(m+\nu)!(m+\nu+2)!} \sum_{j=0}^{\nu} \binom{m+\nu}{\nu-j} \binom{m+\nu+1}{j}}{\sum_{\nu=0}^{\infty} \frac{(-b)^\nu}{(m+\nu+1)!(m+\nu+2)!} \sum_{j=0}^{\nu} \binom{m+\nu+1}{\nu-j} \binom{m+\nu+1}{j}} \\ &= \frac{2}{b} \frac{\sum_{\nu=0}^{\infty} \frac{(-b)^\nu}{(m+\nu)!(m+\nu+2)!} \binom{2m+2\nu+1}{\nu}}{\sum_{\nu=0}^{\infty} \frac{(-b)^\nu}{(m+\nu+1)!(m+\nu+2)!} \binom{2m+2\nu+2}{\nu}} = \frac{1}{b} \frac{\sum_{\nu=0}^{\infty} \frac{(-b)^\nu (2m+\nu+2)}{(m+\nu+1)!(m+\nu+2)!} \binom{2m+2\nu+2}{\nu}}{\sum_{\nu=0}^{\infty} \frac{(-b)^\nu}{(m+\nu+1)!(m+\nu+2)!} \binom{2m+2\nu+2}{\nu}} \\ &= \frac{2}{b} \frac{\sum_{\nu=0}^{\infty} \frac{(-b)^\nu}{(2m+2\nu+2)!} \binom{2m+2\nu+2}{m+\nu} \binom{2m+2\nu+1}{\nu}}{\sum_{\nu=0}^{\infty} \frac{(-b)^\nu}{(2m+2\nu+3)!} \binom{2m+2\nu+3}{m+\nu+1} \binom{2m+2\nu+2}{\nu}} = \frac{2}{b} \frac{\sum_{\nu=0}^{\infty} \frac{(2m+2\nu+1)!}{(m+\nu)!(m+\nu+2)!(2m+\nu+1)!\nu!} (-b)^\nu}{\sum_{\nu=0}^{\infty} \frac{(2m+2\nu+2)!}{(m+\nu+1)!(m+\nu+2)!(2m+\nu+2)!\nu!} (-b)^\nu}. \end{aligned}$$

Unfortunately the series expansions here have large and oscillating terms, so dealing with it does not seem to be simple. When, e.g., b is of the order m^2 , then also the terms with $\nu \approx m$ are the highest, and there are a large number of similar order large terms. Therefore, this series expansion does not seem to be suitable neither for the computation of the value of the ratio, nor for the extraction of a good polynomial approximation which would approach the global maximum while remaining manageable.

6. APPROXIMATE MAXIMIZATION BY POLYNOMIALS

5.1. In the aimed applications in showing small gaps between consecutive prime numbers it is very important to have a suitably nice, manageable function P . It is enough to mention that even in the simplest case of $P(x) = x^{k+\ell}$, $\ell \asymp \sqrt{k}$ the technical difficulties become rather serious when k and ℓ tend to infinity with the size N of the primes, see [5]. The details of these aspects, when the choice of the weight function is done according to the present work, will be handled in the forthcoming paper [8]. Here we will only present the foreseen choice of the weight P , and show its approximate optimality. The said choice will be a relatively simple function, actually a real polynomial $P(x)$, satisfying the conditions

$$(31) \quad x^k | P(x), \quad P(1) > 0, \quad \deg P(x) = k + C_0 k^{1/3},$$

which is essentially optimal in the extremal problem (1). More exactly, with the notations

$$(32) \quad A_k := \int_0^1 \frac{x^{k-2}}{(k-2)!} (P^{(k-1)}(1-x))^2 dx, \quad B_k := \int_0^1 \frac{x^{k-1}}{(k-1)!} (P^{(k)}(1-x))^2 dx$$

it satisfies with an absolute constant C_1

$$(33) \quad A_k(k + C_1 k^{1/3}) - 4B_k \geq 0.$$

Equivalently,

$$(34) \quad S(P, k) := \frac{A_k}{B_k} \geq \frac{4}{k + C_1 k^{1/3}},$$

in full correspondence with (30). Compared with the optimal transformed Bessel function $q_1(x) = x^{-m/2} J_m(\alpha_{m,1} \sqrt{x})$, the difference is only in the value of the constant C_1 .

In order to define our polynomial we put

$$(35) \quad M := \lceil C_1 k^{1/3} / 6 \rceil, \quad g(y) := (y-1)^4 (2-y)^4.$$

Let us remark that the exact choice of $g(y)$ is irrelevant, any positive polynomial or even a function $g \in C^1[1, 2]$ with a zero of order at least 3 at $y = 1$ and $y = 2$ would suffice for our purposes. After this, let

$$(36) \quad P(x) := P_k(x) := \sum_{\substack{\ell=M \\ 2|\ell}}^{2M} g\left(\frac{\ell}{M}\right) \left(\frac{k}{2}\right)^\ell \frac{x^{k+\ell}}{(k+\ell)!}.$$

5.2. In evaluating A_k and B_k we will use the well-known relation (easily obtained by partial integration and induction) for the Euler integral

$$(37) \quad B(m, n) := \int_0^1 x^n (1-x)^m dx = \frac{m!}{(n+1) \cdots (m+n)} \int_0^1 x^{m+n} dx = \frac{n! m!}{(m+n+1)!}.$$

In view of (36)–(37) we have

$$(38) \quad P^{(k-1)}(x) = \sum_{\substack{\ell=M \\ 2 \nmid \ell}}^{2M} g\left(\frac{\ell}{M}\right) \left(\frac{k}{2}\right)^\ell \frac{x^{\ell+1}}{(\ell+1)!},$$

$$(39) \quad P^{(k)}(x) = \sum_{\substack{\ell=M \\ 2 \nmid \ell}}^{2M} g\left(\frac{\ell}{M}\right) \left(\frac{k}{2}\right)^\ell \frac{x^\ell}{\ell!},$$

$$(40) \quad (P^{(k-1)}(x))^2 = \sum_{\substack{\ell_1=M \\ 2 \nmid \ell_1}}^{2M} \sum_{\substack{\ell_2=M \\ 2 \nmid \ell_2}}^{2M} g\left(\frac{\ell_1}{M}\right) g\left(\frac{\ell_2}{M}\right) \left(\frac{k}{2}\right)^{\ell_1+\ell_2} \frac{x^{\ell_1+\ell_2+2}}{(\ell_1+1)!(\ell_2+1)!},$$

$$(41) \quad (P^{(k)}(x))^2 = \sum_{\substack{\ell_1=M \\ 2 \nmid \ell_1}}^{2M} \sum_{\substack{\ell_2=M \\ 2 \nmid \ell_2}}^{2M} g\left(\frac{\ell_1}{M}\right) g\left(\frac{\ell_2}{M}\right) \left(\frac{k}{2}\right)^{\ell_1+\ell_2} \frac{x^{\ell_1+\ell_2}}{\ell_1! \ell_2!},$$

$$(42) \quad A_k = \sum_{\substack{\ell_1=M \\ 2 \nmid \ell_1}}^{2M} \sum_{\substack{\ell_2=M \\ 2 \nmid \ell_2}}^{2M} g\left(\frac{\ell_1}{M}\right) g\left(\frac{\ell_2}{M}\right) \frac{(k/2)^{\ell_1+\ell_2}}{(k+\ell_1+\ell_2+1)!} \binom{\ell_1+\ell_2+2}{\ell_1+1},$$

$$(43) \quad B_k = \sum_{\substack{\ell_1=M \\ 2 \nmid \ell_1}}^{2M} \sum_{\substack{\ell_2=M \\ 2 \nmid \ell_2}}^{2M} g\left(\frac{\ell_1}{M}\right) g\left(\frac{\ell_2}{M}\right) \frac{(k/2)^{\ell_1+\ell_2}}{(k+\ell_1+\ell_2)!} \binom{\ell_1+\ell_2}{\ell_1}.$$

In the following, put

$$(44) \quad u := \ell_1 + 1, \quad v := \ell_2 + 1, \quad \text{and} \quad H := (u + v)/2, \quad D := (u - v)/2.$$

Taking into account $2 \nmid \ell_1, \ell_2$ and $g(1) = g(2) = 0$, the even variables u, v will run from $M + 1$ to $2M$ and we will have

$$(45) \quad \frac{k^{\ell_1+\ell_2} k!}{(k+\ell_1+\ell_2)!} = 1 + O\left(\frac{M^2}{k}\right).$$

Here and elsewhere in the sequel the implied absolute constants of the O symbol as well as the absolute constants C_i ($i \geq 2$) will be always independent from C_1 .

Clearly, we can replace $A'_k := k!A_k$ and $B'_k := k!B_k$ for A_k and B_k , resp., in (33), hence in order to show (33) it suffices to prove

$$(46) \quad \sum_{H=M+1}^{2M} I_k(H) \left(1 + O\left(\frac{M^2}{k}\right)\right) \geq 0,$$

where

$$(47) \quad I_k(H) := 2^{-2H} \sum_{\substack{2|u,v, u+v=2H \\ u,v \in (M, 2M)}} g\left(\frac{u-1}{M}\right) g\left(\frac{v-1}{M}\right) \binom{u+v-2}{u-1} \left\{ \frac{k + C_1 k^{1/3}}{k + 4M} \frac{(u+v)(u+v-1)}{uv} - 4 \right\}.$$

Let us denote the above summation conditions simply by $\sum_{u,v}^*$ and let us consider first

$$(48) \quad \begin{aligned} I'_k(H) &:= 2^{-2H} \sum_{u,v}^* g\left(\frac{u-1}{M}\right) g\left(\frac{v-1}{M}\right) \binom{u+v-2}{u-1} \left\{ \frac{(u+v)(u+v-1)}{uv} - 4 \right\} \\ &= 2^{-2H} \sum_{u,v}^* g\left(\frac{u-1}{M}\right) g\left(\frac{v-1}{M}\right) \binom{u+v-2}{u-1} \frac{4D^2 - 2H}{H^2 - D^2}. \end{aligned}$$

We will see later that the mere significance of the weight function g is to cut the tails when H is near to M or $2M$, so first we will investigate the simpler sum

$$(49) \quad I''_k(H) := 2^{-2H} \sum_{u,v}^* \binom{u+v-2}{u-1} \frac{4D^2 - 2H}{H^2 - D^2},$$

when

$$(50) \quad H \in [M + t(M), 2M - t(M)], \quad t(M) := 4\sqrt{M \log M}.$$

By Stirling's formula

$$(51) \quad \log \Gamma(s) = \left(s - \frac{1}{2}\right) \log s - s + \frac{1}{2} \log(2\pi) + O\left(\frac{1}{|s|}\right),$$

we obtain

$$(52) \quad \begin{aligned} \log \left(2^{-2H} \binom{u+v-2}{u-1} \right) &= \log \left(\frac{\Gamma(2H-1)2^{-2H}}{\Gamma(H+D)\Gamma(H-D)} \right) \\ &= \left(2H - \frac{3}{2}\right) \left(\log H + \log 2 + \log \left(1 - \frac{1}{2H}\right) \right) + 1 - \frac{\log 2\pi}{2} - 2H \log 2 + O\left(\frac{1}{H}\right) \\ &\quad - \left(H + D - \frac{1}{2}\right) \left(\log H + \log \left(1 + \frac{D}{H}\right) \right) - \left(H - D - \frac{1}{2}\right) \left(\log H + \log \left(1 - \frac{D}{H}\right) \right) \\ &= \frac{-\log H}{2} - \frac{3 \log 2}{2} - 1 + 1 - \frac{\log 2\pi}{2} \\ &\quad - \sum_{n=0}^{\infty} \left(\frac{2D}{2n+1} \left(\frac{D}{H}\right)^{2n+1} - \frac{2H}{2n+2} \left(\frac{D}{H}\right)^{2n+2} \right) + O\left(\frac{1}{H} + \frac{D^2}{H^2}\right). \\ &= \frac{-\log H}{2} - \frac{\log 16\pi}{2} - \sum_{n=0}^{\infty} \frac{1}{(n+1)(2n+1)} \frac{D^{2n+2}}{H^{2n+1}} + O\left(\frac{1}{H} + \frac{D^2}{H^2}\right). \end{aligned}$$

Using $e^{-|x|} = 1 + O(x)$ we obtain from (49), (50) and (52)

$$(53) \quad \begin{aligned} I''_k(H) &= \frac{1}{2\sqrt{\pi}H^{3/2}} \sum_{\substack{2|D-H \\ M < H-D, H+D \leq 2M}} e^{-D^2/H} \left(2\frac{D^2}{H} - 1 \right) \left(1 + O\left(\frac{1}{H} + \frac{D^2}{H^2} + \frac{D^4}{H^3}\right) \right) \\ &= \frac{I_k^*(H)}{2\sqrt{\pi}H^{3/2}} + O\left(\frac{1}{M^2}\right) + O\left(\int_{t(M)}^{\infty} \frac{t^2}{H} e^{-t^2/H} dt\right) = \frac{I_k^*(H)}{2\sqrt{\pi}H^{3/2}} + O\left(\frac{1}{M^2}\right), \end{aligned}$$

where

$$(54) \quad I_k^*(H) := \sum_{\substack{2|m-H \\ m=-\infty}}^{\infty} f(m), \quad f(x) = \left(2\frac{x^2}{H} - 1 \right) e^{-x^2/H}.$$

5.3. We will use Poisson summation formula

$$(55) \quad \sum_{n=-\infty}^{\infty} f(t+nT) = \frac{1}{T} \sum_{\nu=-\infty}^{\infty} \widehat{f}\left(\frac{\nu}{T}\right) e^{2\pi i \nu t/T}$$

with $T = 2$, $t = 0$ for $2|H$ and $T = 2$, $t = 1$ for $2 \nmid H$, valid if⁴

$$(56) \quad |f(x)| + |\widehat{f}(x)| \leq C(1 + |x|^{-1-\delta}) \quad \text{with some } \delta > 0, C > 0.$$

Let us consider first the case when H is even, that is when $m = 2n$ in the above sum (54). We have then

$$\begin{aligned} 2I_k^*(H) &= 2 \sum_{n=-\infty}^{\infty} f(2n) = \sum_{\nu=-\infty}^{\infty} \widehat{f}(\nu/2) = \sum_{\nu=-\infty}^{\infty} \int_{-\infty}^{\infty} f(x) e^{\pi i \nu x} dx \\ &= \sum_{\nu=-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-x^2/H} \left(2\frac{x^2}{H} - 1\right) e^{\pi i \nu x} dx = \sqrt{H} \sum_{\nu=-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-y^2} (2y^2 - 1) e^{\pi i \nu \sqrt{H} y} dy \\ &= \sqrt{H} \sum_{\nu=-\infty}^{\infty} \int_{\Im y=0} e^{-(y-\pi i \nu \sqrt{H}/2)^2 - \pi^2 \nu^2 H/4} \\ &\quad \cdot \left\{2(y - \pi i \nu \sqrt{H}/2)^2 + 2(y - \pi i \nu \sqrt{H}/2) \pi i \nu \sqrt{H} - \pi^2 \nu^2 H/2 - 1\right\} dy \\ &= \sqrt{H} \sum_{\nu=-\infty}^{\infty} \int_{\Im z = -\pi \nu \sqrt{H}/2} e^{-z^2 - \pi^2 \nu^2 H/4} \left\{2z^2 + 2\pi i \nu \sqrt{H} z - \pi^2 \nu^2 H/2 - 1\right\} dz \\ &= \sqrt{H} \sum_{\nu=-\infty}^{\infty} \int_{\Im z=0} e^{-z^2 - \pi^2 \nu^2 H/4} \left\{2z^2 + 2\pi i \nu \sqrt{H} z - \pi^2 \nu^2 H/2 - 1\right\} dz \\ (57) \quad &= \sqrt{H} \int_{z \in \mathbb{R}} e^{-z^2} (2z^2 - 1) dz + O(e^{-2H}) = \sqrt{H} \left[-ze^{-z^2}\right]_{-\infty}^{\infty} + O(e^{-2H}) = O(e^{-2H}). \end{aligned}$$

The proof runs completely analogously for $t = 1$, that is, when H is odd, because the extra factor $e^{\pi i \nu}$ does not change the modulus of $\widehat{f}(\nu/2)$.

5.4. Dealing with the original integral $I'_k(H)$ we have to take into account the effect of the weight function g as well. From the Taylor expansion of g we find

$$(58) \quad g\left(\frac{H \pm D - 1}{M}\right) = g\left(\frac{H - 1}{M}\right) \pm g'\left(\frac{H - 1}{M}\right) \frac{D}{M} + O\left(\frac{D^2}{M^2}\right)$$

so this effect is

$$(59) \quad g\left(\frac{H + D - 1}{M}\right) g\left(\frac{H - D - 1}{M}\right) = g^2\left(\frac{H - 1}{M}\right) + O\left(\frac{D^2}{M^2}\right).$$

The effect of the error term on $I'_k(H)$ is here, similarly to (49),

$$(60) \quad O\left(H^{-3/2} \sum_{\substack{2|D-H \\ |D| < M}} e^{-D^2/H} \left(1 + \frac{D^2}{H}\right) \left(1 + \frac{D^4}{H^3}\right) \frac{D^2}{M^2}\right) = O\left(\frac{1}{M^2}\right).$$

⁴For this form see [12, Chapter VII, §2].

So we obtain for all integers H subject to (50) from (48), (49), (53), (55), (57), (59) and (60)

$$(61) \quad I'_k(H) = I''_k(H)g^2\left(\frac{H-1}{M}\right) + O\left(\frac{1}{M^2}\right) = \frac{g^2\left(\frac{H-1}{M}\right)}{2\sqrt{\pi}H^{3/2}}I_k^*(H) + O\left(\frac{1}{M^2}\right) \geq -\frac{C_2}{M^2}.$$

On the other hand, if (50) does not hold, that is if

$$(62) \quad \min(H-M, 2M-H) < t(M),$$

then we find $|D| < t(M)$ and $|H \pm D - 1| \in [M, M+2t(M)] \cup [2M-2t(M), 2M]$, whence

$$(63) \quad g\left(\frac{H+D-1}{M}\right)g\left(\frac{H-D-1}{M}\right) \ll \frac{t^4(M)}{M^4} \ll \frac{1}{M^{7/4}}.$$

This implies in case of (62)

$$(64) \quad I'_k(H) \ll M^{-7/4}M^{-3/2} \int_{-\infty}^{\infty} e^{-u^2/H} \left(\frac{u^2}{H} + 1\right)^3 du \ll M^{-11/4}.$$

Consequently, for $k > k_0$ we have by $I_k(H) \geq I'_k(H)$

$$(65) \quad \sum_{\substack{H \\ \min(H-M, 2M-H) < t(M)}} I_k(H) \geq \sum_{\substack{H \\ \min(H-M, 2M-H) < t(M)}} I'_k(H) \geq -C_3M^{-2}.$$

Finally, let us investigate now for H in (50) the difference

$$(66) \quad I_k(H) - I'_k(H) \geq \frac{M}{k} \cdot 3\tilde{I}_k(H),$$

where similarly to the above considerations

$$(67) \quad \begin{aligned} \tilde{I}_k(H) &:= 2^{-2H} \sum_{u,v}^* g\left(\frac{u-1}{M}\right)g\left(\frac{v-1}{M}\right) \binom{u+v-2}{u-1} \\ &= \frac{1}{2\sqrt{\pi}H} \sum_{\substack{2|D-H \\ M < H-D, H+D \leq 2M}} e^{-D^2/H} \left(1 + O\left(\frac{1}{H} + \frac{D^2}{H^2} + \frac{D^4}{H^3}\right)\right) \\ &\quad \cdot \left(g^2\left(\frac{H-1}{M}\right) + O\left(\frac{D^2}{M^2}\right)\right) \geq C_4g^2\left(\frac{H-1}{M}\right) - \frac{C_5}{M}. \end{aligned}$$

Summing over all H in (62) we obtain

$$(68) \quad \sum_{H=M+t(M)}^{2M-t(M)} I_k(H) \geq \sum_{H=M+t(M)}^{2M-t(M)} I'_k(H) + \frac{C_6M^2}{k}.$$

Therefore by (61) and (65)–(68) we have finally

$$(69) \quad \sum_{H=M+1}^{2M} I_k(H) \geq \frac{C_6M^2}{k} - \frac{C_2}{M} - \frac{C_3}{M^2} > 0,$$

if C_1 was chosen sufficiently large, satisfying

$$(70) \quad C_1 > 6 \left(\frac{C_2}{C_6}\right)^{1/3}.$$

REFERENCES

- [1] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series 55, For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] F. Bowman, *Introduction to Bessel functions*, Dover Publications Inc., New York, 1958.
- [3] J. Friedlander and H. Iwaniec, *Opera de cribro*, American Mathematical Society Colloquium Publications 57, American Mathematical Society, Providence, RI, 2010.
- [4] D. A. Goldston, J. Pintz and C. Y. Yıldırım, Primes in tuples. I, *Ann. of Math. (2)* **170** (2009), 819–862.
- [5] ———, Primes in tuples. II, *Acta Math.* **204** (2010), 1–47.
- [6] D. A. Goldston, Y. Motohashi, J. Pintz and C. Y. Yıldırım, Small gaps between primes exist, *Proc. Japan Acad. Ser. A Math. Sci.* **82** (2006), 61–65.
- [7] P. Meyer-Nieberg, *Banach lattices*, Universitext, Springer-Verlag, Berlin, 1991.
- [8] J. Pintz, Some new results on gaps between consecutive primes, (2013), to appear.
- [9] F. Riesz and B. Sz.-Nagy, *Functional analysis*, Dover Books on Advanced Mathematics, Dover Publications Inc., New York, 1990, Translated from the second French edition by Leo F. Boron, Reprint of the 1955 original.
- [10] K. Soundararajan, Note on Goldston-Pintz-Yıldırım, (an analysis of the GPY result), (2005), unpublished note, sent in an email letter to D. Goldston.
- [11] ———, Small gaps between prime numbers: the work of Goldston-Pintz-Yıldırım, *Bull. Amer. Math. Soc. (N.S.)* **44** (2007), 1–18.
- [12] E. M. Stein and G. Weiss, *Introduction to Fourier analysis on Euclidean spaces*, Princeton University Press, Princeton, N.J., 1971, Princeton Mathematical Series, No. 32.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS
HUNGARIAN ACADEMY OF SCIENCES
1364 BUDAPEST, PO Box 127, HUNGARY
E-mail address: pintz@renyi.mta.hu
revesz.szilard@renyi.mta.hu

DEPARTMENT OF MATHEMATICS
KUWAIT UNIVERSITY
P.O. BOX 5969 SAFAT – 13060 KUWAIT
E-mail address: szilard@sci.kuniv.edu.kw

BERGISCHE UNIVERSITÄT WUPPERTAL
FACULTY OF MATHEMATICS AND NATURAL SCIENCE
GAUSSSTRASSE 20, 42119 WUPPERTAL, GERMANY
E-mail address: farkas@uni-wuppertal.de