

Problems - 2012.10.24

P. Maga & P. P. Pach

1. Let $p_1, \dots, p_k, q_1, \dots, q_l$ be prime numbers. Prove that there is a prime number P such that p_1, \dots, p_k are all quadratic residues and q_1, \dots, q_l are all quadratic nonresidues modulo P .
2. For a given prime p , denote by $l(p)$ the least quadratic nonresidue. Prove that for any $\varepsilon > 0$, $l(p) < \varepsilon p$ if p is large enough.
3. Determine all positive integers relatively prime to all the terms of the infinite sequence

$$a_n = 2^n + 3^n + 6^n - 1, \quad n \geq 1.$$

4. Let t be a fixed integer. Show that for any prime number p , one can find a positive integer n such that

$$(3 - 7t)2^n + (18t - 9)3^n + (6 - 10t)4^n$$

is divisible by p .

5. Determine all positive integers $n \geq 2$ that satisfy the following condition: for all a and b relatively prime to n we have $a \equiv b \pmod{n}$ if and only if $ab \equiv 1 \pmod{n}$.
6. Let p be an odd prime and let \mathbb{Z}_p denote (the field of) integers modulo p . How many elements are in the set

$$\{x^2 : x \in \mathbb{Z}_p\} \cap \{y^2 + 1 : y \in \mathbb{Z}_p\}?$$

7. For which prime p does there exist a cubic polynomial f of integer coefficients such that the leading coefficient is not divisible by p and $f(1), \dots, f(p)$ are distinct modulo p ?