## Problems - 2013.10.31

P. Maga & P. P. Pach

1. Find all differentiable functions  $f: \mathbb{R} \to \mathbb{R}$  such that

$$f'(x) = \frac{f(x+n) - f(x)}{n}$$

for all real numbers x and all positive integers n.

2. Let  $F_0(x) = \ln x$ . For  $n \ge 0$  and x > 0, let  $F_{n+1}(x) = \int_0^x F_n(t) dt$ . Evaluate

$$\lim_{n\to\infty}\frac{n!F_n(1)}{\ln n}.$$

3. Suppose that  $f:[0,1]\to\mathbb{R}$  has a continuous derivative and that  $\int_0^1 f(x) dx = 0$ . Prove that for every  $\alpha\in(0,1)$ ,

$$\left| \int_0^\alpha f(x) \, dx \right| \le \frac{1}{8} \max_{0 \le x \le 1} |f'(x)|.$$

4. Let f be a real function on the real line with continuous third derivative. Prove that there exists a point a such that

$$f(a) \cdot f'(a) \cdot f''(a) \cdot f'''(a) \ge 0.$$

5. Show that for every positive integer n,

$$\left(\frac{2n-1}{e}\right)^{\frac{2n-1}{2}} < 1 \cdot 3 \cdot 5 \cdots (2n-1) < \left(\frac{2n+1}{e}\right)^{\frac{2n+1}{2}}.$$

6. Let  $f: \mathbb{R} \to \mathbb{R}$  be a continuous function. Suppose that for any c > 0, the graph of f can be moved to the graph of cf using only a translation or a rotation. Does this imply that f(x) = ax + b for some real numbers a and b?